

POLYMER CHEMISTRY

AN INTRODUCTION

Raymond B. Seymour
Charles E. Carraher, Jr.

Glossary	76
Exercises	80
Bibliography	82
4 / MOLECULAR WEIGHT OF POLYMERS	87
4.1 Introduction	87
4.2 Average Molecular Weight Values	89
4.3 Fractionation of Polydisperse Systems	93
4.4 Gel Permeation Chromatography (GPC)	94
4.5 End Group Analysis	96
4.6 Ebulliometry and Cryometry	96
4.7 Osmometry	97
4.8 Refractometry	100
4.9 Light-Scattering Measurements	101
4.10 Ultracentrifugation	106
Summary	107
Glossary	108
Exercises	111
Bibliography	113
5 / TESTING AND CHARACTERIZATION OF POLYMERS	122
5.1 Typical Stress-Strain Curves	122
5.2 Physical Tests	124
5.3 Spectronic Characterization of Polymers	129
5.4 Thermal Analysis	137
Summary	141
Glossary	142
Exercises	145
Bibliography	146
6 / NATURALLY OCCURRING POLYMERS	155
6.1 Polysaccharides	156
6.2 The Xanthate Viscose Process	158
6.3 Chitin	159
6.4 Starch	159
6.5 Other Polysaccharides	162
6.6 Proteins	162
6.7 Nucleic Acids	166
6.8 Naturally Occurring Polyisoprenes	170
Summary	179
Glossary	181
Exercises	186
Bibliography	188

7 / STEP-REACTION POLYMERIZATION OR POLYCONDENSATION REACTIONS	193
7.1 Comparison between Polymer Type and Kinetics of Polymerization	193
7.2 Introduction	202
7.3 Stepwise Kinetics	203
7.4 General Step-Reaction Polymerization	211
7.5 Polyesters	213
7.6 Synthetic Polyamides	216
7.7 Polyimides	218
7.8 Polybenzimidazoles and Related Polymers	218
7.9 Polyurethanes and Polyureas	219
7.10 Polysulfides	221
7.11 Polyethers	221
7.12 Polysulfones	223
7.13 Phenolic and Amino Plastics	223
7.14 General Information on Step-Reaction Polymerization	227
7.15 Polycondensation Mechanisms	228
7.16 Synthetic Routes	230
Summary	235
Glossary	237
Exercises	242
Bibliography	244
8 / IONIC CHAIN-REACTION AND COMPLEX COORDINATIVE POLYMERIZATION (ADDITION POLYMERIZATION)	250
8.1 Cationic Polymerization	250
8.2 Anionic Polymerization	258
8.3 Polymerization with Complex Coordination Catalysts	263
8.4 Polymers of 1,3-Butadiene	266
8.5 Stereoregularity	267
Summary	270
Glossary	271
Exercises	274
Appendix	277
Bibliography	277
9 / FREE-RADICAL CHAIN POLYMERIZATION (ADDITION POLYMERIZATION)	281
9.1 Initiators for Free-Radical Chain Polymerization	281
9.2 Mechanism for Free-Radical Chain Polymerization	287
9.3 Chain Transfer	295
9.4 Polymerization Techniques	300

In the mechanism, is widely used for the production of many commercial plastics and elastomers. While the particles in the suspension range from 10 to 1000 nm, those in the emulsion process range from 0.05 to 5 nm. The small beads produced in the suspension process may be separated by filtering, but the latex produced in emulsion polymerization is a stable system in which the charged particles cannot be removed by ordinary separation procedures.


Since relatively stable macroradicals are produced in the emulsion process, the termination rate is decreased and a high molecular weight product is produced at a rapid rate. It is customary to use a water-soluble initiator such as potassium persulfate, and an anionic surfactant such as sodium stearate and to stir the aqueous mixture of monomer, initiator, and surfactant in the absence of oxygen at 40 to 70°C.

A typical recipe for emulsion polymerization includes 100 g of monomer, such as styrene, 180 g of water, 5 g of sodium stearate, and 0.5 g of potassium persulfate. When the concentration of soap exceeds the critical micelle concentration (CMC), these molecules are present as micelles in which the hydrophilic carboxylic acid ends are oriented toward the water-micelle interface, and the lyophobic hydrocarbon ends are oriented toward the center of the micelle. The micelles are present as spheres with a diameter of 5 to 10 nm when the soap concentration is less than 2%. However, with the higher concentrations customarily used, the micelles resemble aggregates of rods which are 100 to 300 nm in length.

As shown in Fig. 9.3, the water-insoluble monomer is attracted to the lyophobic ends in the micelles, and this causes the micelles to swell. The number of swollen micelles per milliliter of water is on the order of 10^{18} . However, at the initial stages of polymerization (phase I), most of the monomer is present as globules which resemble those observed in suspension polymerization.

Since the initiation of polymerization takes place in the aqueous phase, essentially no polymerization takes place in the globules. Thus, they serve primarily as a reservoir of monomer supplied to the micelles to replace the monomer converted to polymer. The number of droplets per milliliter of water is on the order of 10^{11} . Hence, since there are 10 million times as many micelles as droplets, the chance of initiation of monomer in a droplet is very, very small.

As shown in the following equations, the persulfate ion undergoes homolytic cleavage to produce two sulfate ion radicals. These serve as initiators for the few water-soluble monomer molecules present in the aqueous phase.



According to a theory proposed by Harkins and refined by Smith and Ewart, the first stages of propagation in an emulsion system also take place in the aqueous phase to produce a more lyophilic surface-active oligoradical, as shown below in (9.36).




FIGURE 9.3 Micelles swollen with solubilized styrene monomer. (From Introduction to Polymer Chemistry by R. Seymour, McGraw-Hill, New York, 1971. Used with permission of McGraw-Hill Book Company.)